

PLANT PATENT APPLICATION

Strawberry Variety Named 'Albion'

Inventor: Douglas V. Shaw, a citizen of the United States of America, residing at
1002 Stanford Dr., Davis, California 95616

Kirk D. Larson, a citizen of the United States of America, residing at
4 Russell Ct., Irvine, California 92175

Assignee: Regents of the University of California
Office of Technology Transfer
University of California
Office of the President
1111 Franklin Street, 5th Floor
Oakland, CA 94607-5200

Entity: small

Strawberry Variety Named "Albion"

GENUS AND SPECIES

[0001] The strawberry cultivar of this invention is botanically identified as *Fragaria x ananassa* Duch.

VARIETY DENOMINATION

[0002] The variety denomination is 'Albion'.

BACKGROUND OF THE INVENTION

[0003] 'Albion' originated from a cross performed in 1997 between the cultivar "Diamante" (U. S. Pl. Pat. 10,435) and advanced selection Cal 94.16-1. 'Albion' was first fruited at the University of California Wolfskill Experimental Orchard, near Winters, CA in 1998, where it was selected, originally designated Cal 97.117-3, and propagated asexually by runners. Following selection and during testing, the plant was originally designated 'CN220', and subsequently has been named 'Albion' for introduction. Asexual propagules from this original source have been tested at the Watsonville Strawberry Research Facility, the South Coast Research and Extension Center, and to a limited extent in grower fields starting in 1999. The properties of this variety were found to be transmissible by such asexual reproduction.

BRIEF SUMMARY OF THE INVENTION

[0004] This invention relates to a new and distinctive day-neutral type cultivar designated as 'Albion'. 'Albion' is a day-neutral (everbearing) cultivar similar to 'Diamante' (U.S. Pl. Pat. 10,435), but with higher quality fruit, lower cull rate, darker fruit, and substantially better resistance to *Phytophthora cactorum*; it is similar to 'Aromas' (U.S. Pl. Pat. 10,451), but with larger, higher quality, firmer and better-flavored fruit.

BRIEF DESCRIPTION OF THE DRAWINGS

[0005] The figures depict various characteristics of the 'Albion' cultivar.

[0006] Figure 1 shows the general flowering and fruiting characteristics of plants in a field planting.

[0007] Figure 2 shows rows of typical fruiting plants.

[0008] Figure 3 shows a typical leaf at mid-season.

[0009] Figure 4 shows representative mid-season fruit.

DETAILED DESCRIPTION OF THE INVENTION

[0010] This invention relates to a new and distinctive day-neutral type cultivar designated as 'Albion'. 'Albion' is typical of day-neutral strawberry cultivars and produces fruit regardless of day length when treated appropriately in arid, subtropical climates. 'Albion' is moderate to weak in expressing the day-neutral character, being comparable or slightly more day-neutral than 'Diamante' (U. S. Pl. Pat. 10,435), and less so than 'Fern' (U. S. Pl. Pat. 5267) or 'Irvine' (U. S. Pl. Pat. 7172). The production pattern for 'Albion' is similar to that for 'Diamante', although it is somewhat later to reach peak fruiting with most cultural treatments. 'Albion' will be of special interest for winter plantings and in summer plantings where 'Diamante' and 'Aromas' have been successful.

[0011] 'Albion' has the following characteristics relative to parent 94.16-1. The 94.16-1 cultivar is a short-day or June-bearing genotype, whereas Albion is day-neutral. Albion has larger and more uniform sized fruit than 94.16-1, its fruit is substantially firmer, and better flavored. The fruit shape of 94.16-1 tends towards a short but symmetrical conic shape; Albion fruit has a long conical shape. Albion has substantially more even external fruit color, whereas 94.16-1 tends to be mottled during some portions of the season.

[0012] Plants and foliage: Fruiting plants of 'Albion' are similar in morphology to 'Diamante' although slightly more erect and more open; 'Albion' plants are substantially more open and erect than 'Aromas' plants. Comparative statistics for foliar characters near mid-season are given for 'Albion' and the two comparison cultivars in Table 1. Individual leaflets for 'Albion' are smaller than for 'Aromas' or 'Diamante', and somewhat less rounded than for 'Diamante'. Leaves (including petioles) for 'Albion' are shorter than those for the comparison cultivars, mostly due to shorter petiole length. Petioles are generally thicker than those of the comparison cultivars and tend to have heavy pubescence. The adaxial (upper) and abaxial (lower) surfaces of leaves for 'Albion' are similar in color to 'Aromas' and 'Diamante' leaves at mid season, but tend to remain darker and less yellow than those of either comparison cultivar late in the harvest season. Leaves of 'Albion' have consistently less concavity than 'Aromas', and are similar in form to 'Diamante', with more, and usually more rounded serrations than the comparison cultivars.

Table 1. Foliar characteristics for 'Albion', 'Aromas', and 'Diamante'.

Foliar Character	Cultivar		
	'Aromas'	'Diamante'	'Albion'
Plant height (mm)			
mean	257	231	252
range	230-330	160-275	210-270
Plant spread (mm)			
mean	318	357	341
range	241-382	292-419	304-394
Mid-tier leaflet			
Length (mm)			
mean	92	77	73
range	67-100	55-110	50-95
Width (mm)			
mean	74	81	68
range	65-85	55-110	50-95
Mid-tier leaf			
Length (mm)			
mean	253	215	185
range	172-305	169-290	153-223
Width (mm)			
mean	156	149	135
range	135-200	90-210	105-170
Leaf components			
Petiole length (mm)			
mean	154	126	105
range	100-225	92-170	70-130
Petiole diameter (mm)			
mean	3.6	3.7	4.1
range	3.0-4.2	3.0-4.8	3.7-4.6
Petiolule length (mm)			
mean	7.2	11.4	7.4
range	7-12	8-15	5-10
# leaflets/leaf	3	3	3
Leaf convexity	some flat, most slight concave	some flat, most slight concave	some flat, most slight concave
Serrations			
number/leaf	65.1	63.1	71.8
range	57-71	45-72	55-87
shape	semi-pointed	semi-pointed	semi-pointed

Foliar Character	Cultivar		
	'Aromas'	'Diamante'	'Albion'
Leaf pubescence	moderate	very light	light-moderate
Petiole pubescence density direction	moderate perpendicular	moderate to heavy perpendicular to slightly acropetal	heavy perpendicular
Petiole color (Munsell)	5 GY 6/8	5 GY 6/8 5 GY 7/10	5 GY 7/10
Stipule length (mm) mean range	26.9 25-31	26.5 20-35	23.3 14-34
Stipule color core margins	5 GY 6/8 2.5 R 7/8	5 GY 6/8 2.5 R 6/11	5 GY 6/8 2.5 R 7/8
Stolons per nursery mother plant	30.3	24.4	22.5
Venation pattern color	pinnate 2.5 GY 6/8	pinnate 2.5 GY 6/8	pinnate 2.5 GY 6/8

[0013] Flowering, fruiting, fruit, and production characteristics: 'Albion' is similar to other California day-neutral cultivars (e.g., 'Diamante' and 'Aromas') in that it will flower independently of day length, given appropriate temperature and horticultural conditions.

- 5 Comparative statistics for flower and fruit characters near mid-season are given for the three cultivars in Table 2. The primary flowers for 'Albion' are similar in size to 'Diamante' and larger than those of 'Aromas'; the sepals are similar in length to both comparison cultivars, but intermediate in width. The calyx for 'Albion' varies in position but is more frequently slightly necked than either comparison cultivar; each primary flower has 5-8 petals.
- 10 Table 2. Flower and fruit characters for 'Albion', 'Diamante', and 'Aromas'.

Character	Cultivar		
	'Aromas'	'Diamante'	'Albion'
Petal number mean range	5.7 5-7	5.4 5-6	

Character	Cultivar		
	'Aromas'	'Diamante'	'Albion'
Petal shape			
apex	truncate to slightly obtuse	truncate to slightly obtuse	truncate to slightly obtuse
base margin	attenuate entire	attenuate entire	attenuate entire
Petal length (mm)			
mean	13.8	13.4	12.7
range	12-16	11-17	11-15
Petal width (mm)			
mean	13.6	13.2	12.6
range	11-16	12-15	11-14
Flower position (relative to foliage)	most even some internal	most even some internal and exposed	most exposed, some even
Calyx diam.(mm)			
mean	28.6	35.2	35.8
range	27-30	24-40	30-39
Corolla diam.(mm)			
mean	26.3	28.8	27.0
range	23-28	21-34	25-30
Sepal length (mm)			
mean	15.8	19.1	18.5
range	15-19	13-25	14-24
Sepal width (mm)			
mean	5.8	9.5	7.1
range	4-8	7-16	5-9
Sepal color (Munsell)	5 GY 5/6	5 GY 5/6	7.5 GY 4/4
Pedicel length (mm)			
mean	129.5	148.2	113.0
range	110-160	110-180	83-190
Pedicel diameter (mm)			
mean	2.4	2.3	2.9
range	1.9-3.3	1.8-2.7	2.2-3.5
Pedicel color	5 GY 6/8	2.5 GY 6/8	5 GY 6/8
Fruit shape			
Fruit length (mm)			
mean	48.4	53.5	60.6
range	45-57	45-60	55-75
Fruit width (mm)			
mean	43.7	51.1	49.7

Character	Cultivar		
	'Aromas'	'Diamante'	'Albion'
range	35-50	45-60	45-55
Length/ width			
ratio	1.1	1.0	1.2
range	0.9-1.3	0.9-1.3	1.1-1.4
subjective	mostly medium to short rounded conic	rounded to flat conic	most long symmetrical conic
Calyx position	even to indented	even to indented	even to slight neck
Seed position	even to indented	mostly indented, some even	mostly indented, some even

[0014] The fruit shape for 'Albion' can vary but is typically a long and symmetrical conic, and is easily distinguished from 'Aromas' (shortened and rounded conic) or 'Diamante' (rounded and occasionally flattened conic); 'Albion' usually has a greater proportion of symmetrical fruit than either comparison cultivar. External and internal fruit color for 'Albion' is darker than for 'Diamante' and slightly lighter than for 'Aromas' with substantially greater red color (Table 3). Achenes vary from yellow to dark red, and are even with the fruit surface or slightly indented.

Table 3. Foliar and fruit color characteristics for 'Albion', 'Aromas', and 'Diamante'.

Color Character	Cultivar		
	'Aromas'	'Diamante'	'Albion'
Leaf color (CIELAB)			
Adaxial			
L*			
mean	32.7	32.4	32.7
range	31.1-34.2	29.9-35.8	31.5-34.1
a*			
mean	-7.0	-8.1	-7.4
range	-5.8 - -8.2	-7.3 - -10.2	-6.3 - -8.1
b*			
mean	11.8	12.0	11.7
range	9.6-14.0	9.1-13.9	10.4-13.1
Munsell	2.5 GY 3/3	10GY 3/2	5GY 3/2
Abaxial			
L*			
mean	52.3	50.7	49.1
range	34.1-52.3	48.3-52.4	48.6-52.1
a*			

Color Character	Cultivar		
	'Aromas'	'Diamante'	'Albion'
mean	-8.3	-9.0	-8.8
range	-6.8 - -8.8	-8.4 - -9.5	-7.6 - -9.2
b*			
mean	18.1	19.1	19.3
range	13.1-20.1	17.9-21.3	15.7-21.6
Munsell	5GY 5/6 7.5 GY 5/7	5GY 5/6	5GY 5/6
Fruit color (CIELAB)			
External			
L*			
mean	38.1	43.7	40.0
range	36.6-41.4	39.5-47.4	34.3-44.8
a*			
mean	39.3	41.7	41.2
range	37.3-41.2	35.0-46.3	34.8-44.2
b*			
mean	27.2	32.1	28.4
range	21.7-32.9	27.0-35.9	20.9-36.9
Munsell	5 R 3/7	7.5 R 4/11	5R 3/7
Internal			
L*			
mean	65.1	68.3	63.2
range	56.4-70.6	63.7-71.1	56.3-64.4
a*			
mean	31.2	23.9	31.1
range	13.3-40.3	15.5-30.6	20.1-35.9
b*			
mean	34.5	29.2	33.1
range	17.1-41.7	20.3-35.9	24.0-35.9
Munsell	5 R 4/12	7.5 R 6/12	7.5 R 4/11

*CIELAB is the abbreviation of the international color system known as "Commission Internationale De L' Eclairage" 1978. For recommendations concerning uniform color spaces, color difference equations, and psychometric color terms see Supplement No. 2 of CIE Publication No. 15, Paris.

5

[0015] 'Albion' has been tested under a variety of cultural regimes, and optimal performance is obtained when nursery treatments and nutritional programs similar to those for 'Diamante' are used. In general, 'Albion' is very similar in vigor to 'Diamante' and requires less chilling to maintain excellent fruit quality than 'Aromas'. 'Albion' retains good fruit quality in summer planting systems, similar to 'Diamante'.

10

[0016] It is possible that the phenotype may vary somewhat with variations in the environment. Phenotypic features may also vary depending on culture conditions.

[0017] When treated with appropriate planting regimes, 'Albion' has similar fruit size and produces similar individual-plant yields to 'Diamante'; it produces less per plant but develops higher quality fruit than 'Aromas' (Table 4). 'Albion' has a similar production pattern to 'Diamante', although the production is less peaked and less affected by yearly variation in climate. Commercial appearance ratings have been better than those for 'Diamante' and substantially better than those for 'Aromas'; these superior appearance scores translate directly into a smaller fraction of non-marketable fruit than is produced by the comparison cultivars. Fruit for 'Albion' is substantially firmer than fruit from 'Aromas', slightly less firm than 'Diamante'. Subjectively, 'Albion' has outstanding flavor. The fruit will be exceptional for both fresh market and processing, and will be useful for home garden purposes.

Table 4. Performance of 'Albion', 'Aromas', and 'Diamante' evaluated at the Watsonville Research Facility in 2001 and 2002. All plants for these trials were harvested from Macdoel on October 15, and transplanted after 20-28 days supplemental storage. Harvest was initiated in early April and continued through the last week of October. Late yield is that accumulated after August 15 (52" 2-row beds, 17,300 plants/acre).

Item	Yield (g/plant)	Late Yield (g/plant)	Appearance Score (5=best)	Fruit Size (g/fruit)	Firmness
'Aromas'	2,762	582	3.3	28.0	8.6
'Diamante'	2,346	456	3.6	33.5	9.8
'Albion'	2,417	522	4.0	33.0	9.3

[0018] Disease and pest reaction: 'Albion' is moderately resistant to common leaf spot (*Ramularia tulasnei*) and powdery mildew (*Sphaerotheca macularis*). It is quite resistant to Verticillium wilt (*Verticillium dahliae*) and *Phytophthora* crown rot (*Phytophthora cactorum*), and moderately resistant to Anthracnose crown rot (*Colletotrichum acutatum*) (Table 5). When treated properly, it has tolerance to two-spotted spider mites (*Tetranychus*

urticae) equal to that for ‘Diamante’ and better than that for ‘Aromas’. ‘Albion’ is tolerant to strawberry viruses encountered in California.

Table 5. Disease resistance scores for ‘Albion’, ‘Aromas’, and ‘Diamante’; *Phytophthora* and *Verticillium* scores were obtained in evaluations conducted in 2000-2003, *Colletotrichum* was evaluated in 2003.

Genotype	<i>Phytophthora</i>	<i>Verticillium</i>	<i>Colletotrichum</i>
	Resistance	Resistance	Resistance
	Score	Score	Score
	(5=best)	(5=best)	(5=best)
‘Aromas’	4.2	3.5	2.7
‘Diamante’	2.4	2.7	2.6
‘Albion’	4.9	3.4	3.1